

WO 2005/014529

PCT/SE2004/001154

1

2-adamantyl derivatives as P2X₇ receptor antagonists.


The present invention relates to certain adamantane derivatives, processes for their preparation, pharmaceutical compositions containing them, and their use in therapy. In particular, the present invention relates to 2-adamantyl derivatives effective as P2X₇ receptor antagonists.

The P2X₇ receptor (previously known as P2Z receptor), which is a ligand-gated ion channel, is present on a variety of cell types, largely those known to be involved in the inflammatory/immune process, specifically, macrophages, mast cells and lymphocytes (T and B). Activation of the P2X₇ receptor by extracellular nucleotides, in particular adenosine triphosphate, leads to the release of interleukin-1 β (IL-1 β) and giant cell formation (macrophages/microglial cells), degranulation (mast cells) and proliferation (T cells), apoptosis and L-selectin shedding (lymphocytes). P2X₇ receptors are also located on antigen-presenting cells (APC), keratinocytes, salivary acinar cells (parotid cells), hepatocytes and mesangial cells.

It would be desirable to make compounds effective as P2X₇ receptor antagonists for use in the treatment of inflammatory, immune or cardiovascular diseases, in the aetiologies of which the P2X₇ receptor may play a role.

1-Adamantyl derivatives having antagonist activity at the P2X₇ receptor are known in the art and are described, for example, in WO 99/29660, WO 00/61569 and WO 01/94338.

25 The present invention provides a compound of formula (I)



or a pharmaceutically acceptable salt, prodrug or solvate thereof, wherein

A represents a phenyl, pyridyl, indolyl, indazolyl, purinyl, pyrimidinyl, thiophenyl, benzothiazolyl, quinolinyl or isoquinolinyl group, each of which may be optionally substituted by one or more substituents, which may be the same or different, selected from halogen, amino, nitro, cyano, hydroxyl, C₁-C₆ alkyl optionally substituted by at least one substituent selected from hydroxyl or halogen, C₁-C₆ alkoxy, or a group of formula




where Y represents an oxygen or sulphur atom or a group -N(R³)-;

p is 0 or 1;

R¹ represents a bond or a C₁-C₆ alkyl group which may be optionally substituted by at least one substituent selected from hydroxyl, halogen, C₁-C₆ alkoxy, C₁-C₆ alkylthio, C₁-C₆ hydroxyalkyl, C₁-C₆ hydroxyalkoxy, C₁-C₆ alkoxycarbonyl, C₃-C₈ cycloalkyl, phenyl (optionally substituted by at least one substituent selected from halogen, hydroxyl and C₁-C₆ alkylsulphonylamino), benzyl, indolyl (optionally substituted by at least one substituent selected from C₁-C₆ alkoxy), oxopyrrolidinyl, phenoxy, benzodioxolyl, phenoxyphenyl, piperidinyl and benzyloxy;

R² represents hydrogen, hydroxyl, -NR⁴R⁵, except that when R¹ represents a bond, then R² represents a saturated or unsaturated 3- to 10-membered ring system which may comprise at least one ring heteroatom selected from nitrogen, oxygen and sulphur, the ring system being optionally substituted by at least one substituent selected from hydroxyl, amino (-NH₂), C₁-C₆ alkyl, C₁-C₆ alkylamino, -NH(CH₂)₂OH, -NH(CH₂)₃OH, NH(CH₂)₄OH,

C₁-C₆ hydroxyalkyl, benzyl, and



R³ represents a hydrogen atom or a C₁-C₆ alkyl group which may be optionally substituted by at least one substituent selected from hydroxyl, halogen and C₁-C₆ alkoxy;

R⁴ and R⁵ each independently represent hydrogen, pyrrolidinyl, piperidinyl, C₁-C₆ alkylcarbonyl, C₂-C₇ alkenyl, or C₁-C₇ alkyl optionally substituted with at least one substituent selected from carboxyl, hydroxyl, amino (-NH₂), C₁-C₆ alkylamino, di-C₁-C₆

- alkylamino, $-\text{NH}(\text{CH}_2)_2\text{OH}$, $\text{C}_1\text{-C}_6$ alkoxy, $\text{C}_1\text{-C}_6$ alkylthio, $\text{C}_1\text{-C}_6$ alkoxy carbonyl, and a saturated or unsaturated 3- to 10-membered ring system which may comprise at least one ring heteroatom selected from nitrogen, oxygen and sulphur, the ring system being optionally substituted by at least one substituent selected from halogen, hydroxyl, oxo, carboxyl, cyano, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ hydroxyalkyl, $-\text{NR}^6\text{R}^7$, $-(\text{CH}_2)_r\text{NR}^8\text{R}^9$ and $-\text{CONR}^{10}\text{R}^{11}$,
 5 or R^4 and R^5 may together with the nitrogen atom to which they are attached form a saturated 4- to 8-membered heterocyclic ring which may comprise a second ring heteroatom selected from nitrogen and oxygen, the ring being optionally substituted by at least one substituent selected from hydroxyl, halogen, $\text{C}_1\text{-C}_6$ alkyl, and $\text{C}_1\text{-C}_6$ hydroxyalkyl
 10 r is 1, 2, 3, 4, 5 or 6;
 R^6 and R^7 each independently represent a hydrogen atom or a $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_2\text{-C}_6$ hydroxyalkyl or $\text{C}_3\text{-C}_8$ cycloalkyl group, or R^6 and R^7 together with the nitrogen atom to which they are attached form a 3- to 8-membered saturated heterocyclic ring;
 15 R^8 and R^9 each independently represent a hydrogen atom or a $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_2\text{-C}_6$ hydroxyalkyl or $\text{C}_3\text{-C}_8$ cycloalkyl group, or R^8 and R^9 together with the nitrogen atom to which they are attached form a 3- to 8-membered saturated heterocyclic ring; and
 R^{10} and R^{11} each independently represent a hydrogen atom or a $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_2\text{-C}_6$ hydroxyalkyl or $\text{C}_3\text{-C}_8$ cycloalkyl group, or R^{10} and R^{11} together with the nitrogen
 20 atom to which they are attached form a 3- to 8-membered saturated heterocyclic ring;
 B represents $\text{C}(\text{O})\text{NH}$ or $\text{NHC}(\text{O})$;
 n is 1, 2, 3, 4, 5 or 6;
each X is independently selected from halogen or $\text{C}_1\text{-C}_6$ alkoxy; and
 m is 0, 1, 2, 3, 4, 5, 6, 7, 8, or 9;
 25 with the proviso that when B represents $\text{C}(\text{O})\text{NH}$, n is 1 and m is 0, then A is not an unsubstituted phenyl group.

As used herein, the term "halogen" includes fluorine, chlorine, bromine and iodine, and in particular is fluorine or chlorine.

Unless otherwise indicated, the term 'alkyl', when used alone or in combination, refers to a straight or branched chain alkyl moiety. A C₁-C₆ alkyl group has from one to six carbon atoms, including methyl, ethyl, n-propyl, isopropyl, butyl, isobutyl, tert-butyl, n-pentyl, n-hexyl and the like. 'C₁-C₅ alkyl' and 'C₁-C₇ alkyl' will be understood accordingly to mean a straight or branched chain alkyl moiety having from one to five or one to seven carbon atoms respectively.

The term 'cycloalkyl' refers to a saturated alicyclic moiety having from three to eight carbon atoms and includes, for example, cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.

The expression "heterocyclic" includes saturated and unsaturated rings having from 3 up to 10 atoms, at least one of which is a heteroatom selected from oxygen, sulphur or nitrogen. The rings may be mono- or bicyclic and may have alicyclic or aromatic properties. Bicyclic rings may be fully or partially aromatic in character. An unsaturated ring system may be fully or partially unsaturated. Nitrogen heteroatoms will be substituted as necessary, and may also be in the form of *N*-oxides. Sulphur atoms may be in the form of S, S(O) or S(O₂). Where it is intended that the heterocyclic ring has a maximum number of ring atoms that is less than ten, this is specified. Where it is intended that a ring heteroatom is one of N, S or O in particular, or that the heterocyclic ring comprises one or more ring heteroatoms in specific combination, this is indicated.

'Optionally substituted' is used herein to indicate optional substitution by the group or groups specified at any suitable available position.

A "hydroxyalkyl" substituent may contain one or more hydroxyl groups but preferably contains one hydroxyl group.

It will be understood that where B represents C(O)NH the compounds of formula (I) comprise a group A-C(O)NH-(CH₂)_n-; and when B represents NHC(O) the compounds of formula (I) comprise a group A-NHC(O)-(CH₂)_n-.

- 5 In an embodiment of the invention, A represents a phenyl, pyridyl, indolyl or quinolyl group which is optionally substituted as defined above. Preferably, A represents a phenyl, pyridyl or quinolyl group which is optionally substituted as defined above.

- Preferred substituents for the ring A include halogen (for example fluorine or chlorine),
 10 amino, nitro, cyano, hydroxyl, C₁-C₆ alkyl (for example, methyl, ethyl, propyl, butyl, pentyl or hexyl) optionally substituted by at least one substituent selected from hydroxyl and halogen, C₁-C₆ alkoxy (for example methoxy, ethoxy, propoxy, butoxy, pentoxy or hexoxy), or a group of formula -[Y]_p-R¹-R² (II).

- 15 Where Y represents -N(R³)-, R³ is a hydrogen atom or a C₁-C₅ alkyl group which may optionally substituted by at least one substituent (for example, one, two or three substituents independently) selected from hydroxyl, halogen (for example, fluorine, chlorine, bromine or iodine) and C₁-C₆ alkoxy (for example, methoxy or ethoxy).


- 20 Preferably, R³ represents a hydrogen atom or a C₁-C₅ alkyl group which may be optionally substituted by at least one hydroxyl group.

- R¹ represents a bond or a C₁-C₆, preferably C₁-C₄ alkyl group, substituted by at least one
 25 substituent (for example one, two or three substituents independently) selected from hydroxyl, halogen (for example, fluorine or chlorine), C₁-C₆ alkoxy (for example, methoxy or ethoxy), C₁-C₆ alkylthio (for example, methyl- or ethyl-thio), C₁-C₆ hydroxyalkyl (such as hydroxymethyl), C₁-C₆ hydroxyalkyloxy, C₁-C₆ alkoxycarbonyl (for example, methoxycarbonyl), C₃-C₈ cycloalkyl (for example, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl), phenyl (optionally substituted by at least one substituent selected from

halogen, hydroxyl and C₁-C₆ alkylsulphonylamino, such as methylsulphonylamino), benzyl, indolyl (optionally substituted by at least one substituent selected from C₁-C₆ alkoxy such as methoxy), oxopyrrolidinyl, phenoxy, benzodioxolyl, phenoxyphenyl, piperidinyl and benzyloxy.

In an embodiment of the invention, R¹ represents a bond or a C₁-C₄ alkyl group which may be optionally substituted by one, two or three substituents independently selected from hydroxyl, C₁-C₂ alkoxy, methylthio, C₁-C₂ hydroxyalkyl, C₁-C₂ hydroxyalkyloxy, methoxycarbonyl, C₃-C₆ cycloalkyl, phenyl (optionally substituted by at least one substituent selected from halogen, hydroxyl and methylsulphonylamino), benzyl, indolyl (optionally substituted by at least one methoxy), oxopyrrolidinyl, phenoxy, benzodioxolyl, phenoxyphenyl, piperidinyl and benzyloxy.

R² represents hydrogen, hydroxyl, or a group -NR⁴R⁵ except that when R¹ represents a bond, then R² represents a saturated or unsaturated 3- to 10-membered ring system which may comprise at least one ring heteroatom (for example, one, two, three or four ring heteroatoms independently) selected from nitrogen, oxygen and sulphur, the ring system being optionally substituted by at least one substituent (for example, one, two, three or four substituents independently) selected from hydroxyl, amino (-NH₂), C₁-C₆ alkyl, C₁-C₆ alkylamino, -NH(CH₂)₂OH, -NH(CH₂)₃OH, C₁-C₆ hydroxyalkyl, benzyl, and



When R² represents a saturated or unsaturated 3 to 10 membered ring system, this ring system is preferably cyclobutyl, cyclohexyl, bicyclo[2.2.1]hept-2-yl, 2,3-dihydro-1H-indenyl, pyrrolidinyl, piperidinyl or piperazinyl.

In an embodiment of the invention, R^4 and R^5 each independently represent hydrogen, pyrrolidinyl, C_1 - C_2 alkylcarbonyl, C_5 - C_7 alkenyl, or C_1 - C_7 alkyl optionally substituted with one or two substituents independently selected from carboxyl, hydroxyl, amino, C_1 - C_2 alkylamino, di- C_1 - C_2 alkylamino, $-NH(CH_2)_2OH$, C_1 - C_2 alkoxy, C_1 - C_2 alkylthio, C_1 - C_2 alkoxy carbonyl, and a saturated or unsaturated 3- to 10-membered ring system which may comprise at least one ring heteroatom (e.g. one, two, three or four ring heteroatoms independently) selected from nitrogen, oxygen and sulphur, the ring system being optionally substituted by at least one substituent (e.g. one, two, three or four substituents independently) selected from fluorine, hydroxyl, oxo, carboxyl, cyano, C_1 - C_2 alkyl, C_1 - C_2 hydroxyalkyl, $-NR^6R^7$, $-(CH_2)_rNR^8R^9$ and $-CONR^{10}R^{11}$.

The saturated or unsaturated 3- to 10-membered ring system referred to in the preceding paragraph is preferably selected from cyclopropyl, cyclohexenyl, phenyl, thienyl, pyridinyl, furyl, bicyclo[2.2.1]hept-5-en-2-yl, 3,4-dihydro-2H-pyran, thiazolyl, pyrrolyl, pyrazolyl, imidazolyl and thiadiazolyl.

In another embodiment of the invention, where R^4 and R^5 together with the nitrogen atom to which they are attached form a saturated six-membered heterocyclic ring, this ring is optionally substituted by at least one substituent (for example, one, two, three or four substituents independently) selected from hydroxyl, halogen (for example, fluorine, chlorine, bromine or iodine), C_1 - C_6 alkyl (such as methyl or ethyl), and C_1 - C_6 hydroxyalkyl (for example, hydroxymethyl or hydroxyethyl). Preferably, the heterocyclic ring is a piperidinyl, piperazinyl or morpholinyl ring.

R^6 and R^7 each independently represent a hydrogen atom, a C_1 - C_6 alkyl group such as methyl or ethyl, a C_2 - C_6 hydroxyalkyl group (for example, hydroxymethyl) or a C_3 - C_8 cycloalkyl group (such as cyclopentyl or cyclohexyl), or R^6 and R^7 together with the nitrogen atom to which they are attached form a 3 to 8 membered saturated heterocyclic ring (such as pyrrolidinyl or piperidinyl).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35

R⁸ and R⁹ each independently represent a hydrogen atom, a C₁-C₆ alkyl group such as methyl or ethyl, a C₂-C₆ hydroxyalkyl group (for example, hydroxymethyl) or a C₃-C₈ cycloalkyl group (such as cyclopentyl or cyclohexyl), or R⁸ and R⁹ together with the nitrogen atom to which they are attached form a 3 to 8 membered saturated heterocyclic ring (such as pyrrolidinyl or piperidinyl).

R¹⁰ and R¹¹ each independently represent hydrogen, a C₁-C₆ alkyl group such as methyl or ethyl, a C₂-C₆ hydroxyalkyl group (for example, hydroxymethyl) or a C₃-C₈ cycloalkyl group (such as cyclopentyl or cyclohexyl), or R₁₀ and R₁₁ together with the nitrogen atom to which they are attached form a 3 to 8 membered saturated heterocyclic ring (such as pyrrolidinyl or piperidinyl).

Preferably, ring A is unsubstituted or substituted by one or more substituents, which may be the same or different, selected from C₁-C₆ alkyl, optionally substituted by at least one substituent selected from halogen or hydroxyl, or C₁-C₆ alkoxy. Where ring A is substituted, it is preferably substituted by one or two substituents, which may be the same or different.

Preferably n is 1 or 2, especially 1.

X is preferably halogen (such as fluorine, chlorine or bromine) or a C₁-C₆ alkoxy group such methoxy or ethoxy. Most preferably, X is halogen or a C₁-C₄ alkoxy group, especially methoxy.

In an embodiment of the invention, m is 1, 2 or 3.

In another embodiment, m is 0.

It will be appreciated that the number and nature of the substituents on rings in the compounds of the invention will be selected so as to avoid sterically undesirable combinations.

- 5 In one preferred group of compounds according to the invention, A is phenyl, pyridyl or quinolinyl, optionally substituted by one or more substituents, which may be the same or different, selected from C₁-C₆ alkoxy or C₁-C₆ alkyl, optionally substituted by at least one substituent selected from halogen or hydroxyl; B is NHC(O); m is 0; and n is 1

10

Preferred compounds of the invention include:-

2-(2-Adamantyl)-N-(5-methoxy-2-methylphenyl)acetamide,

2-(2-Adamantyl)-N-1*H*-indol-4-ylacetamide,

- 15 2-(1-Adamantyl)-N-quinolin-5-ylacetamide,

or a pharmaceutically acceptable salt, prodrug or solvate thereof.

- Where the compounds according to the invention contain one or more asymmetrically substituted carbon atoms, the invention includes all stereoisomers, including enantiomers
20 and diastereomers, and mixtures including racemic mixtures thereof. Tautomers and mixtures thereof are also included.

- Racemates may be separated into individual enantiomers using known procedures (cf. Advanced Organic Chemistry: 3rd Edition: author J March, p104-107). A suitable
25 procedure involves formation of diastereomeric derivatives by reaction of the racemic material with a chiral auxiliary, followed by separation, for example by chromatography, of the diastereomers and then cleavage of the auxiliary species.


- The compounds according to the invention may be provided as pharmaceutically acceptable
30 salts. Suitable pharmaceutically acceptable salts include base salts such as an alkali metal

salt for example sodium, an alkaline earth metal salt for example calcium or magnesium, an organic amine salt for example triethylamine, morpholine, *N*-methylpiperidine, *N*-ethylpiperidine, procaine, dibenzylamine, *N,N*-dibenzylethylamine or amino acids for example lysine. In another aspect, where the compound is sufficiently basic, suitable salts include acid addition salts such as methanesulphonate, fumarate, hydrochloride, hydrobromide, citrate, maleate and salts formed with phosphoric and sulphuric acid.

Suitable prodrugs of compounds of formula (I) are compounds which are hydrolysed *in vivo* to form compounds of formula (I). These may be prepared by conventional methods.


The present invention further provides a process for the preparation of a compound of formula (I) as defined above, or a pharmaceutically acceptable salt, prodrug or solvate thereof, which comprises:

- (a) when B represents NHC(O) , reacting a compound of formula (III)



wherein L^1 represents a leaving group (e.g. hydroxyl or halogen) and n, m and X are as defined in formula (I), with a compound of formula (IV), A-NH_2 , wherein A is as defined in formula (I); or

- (b) when B represents C(O)NH , reacting a compound of formula



wherein X, m and n are as defined in formula (I), with a compound of formula (VI),
A-C(O)-L², wherein L² represents a leaving group (e.g. hydroxyl or halogen) and A is as
defined in formula (I); and thereafter, if necessary:

converting the compound obtained into a further compound according to the invention
5 and/or forming a pharmaceutically acceptable salt or prodrug or solvate of the compound.

In processes (a) and (b) the coupling reaction is conveniently carried out in an organic
solvent such as dichloromethane, *N,N*-dimethylformamide or 1-methyl-2-pyrrolidinone.

If L¹ or L² represents a hydroxyl group, it may be necessary or desirable to use a coupling
10 agent such as bromo-tris-pyrrolidino-phosphonium hexafluorophosphate (PyBroP).

Compounds of formulae (III) and (V) are known compounds or may be prepared using
known techniques by methods analogous to those known in the art. Examples of
preparation methods for certain of these compounds are given hereinafter in the examples

15

Compounds of formula (I) can be converted into further compounds of formula (I) using
standard procedures conventional in the art.

It will be appreciated that the preparation of compounds of formula (I) may involve, at
20 various stages, the addition and removal of one or more protecting groups. The protection
and deprotection of functional groups is described in 'Protective Groups in Organic
Synthesis', 2nd edition, T.W. Greene and P.G.M. Wuts, Wiley-Interscience (1991).

The compounds of the invention possess activity as P2X₇ receptor antagonists and are
25 therefore indicated as pharmaceuticals for use in the treatment of rheumatoid arthritis,
osteoarthritis, psoriasis, allergic dermatitis, asthma, chronic obstructive pulmonary disease
(COPD), hyperresponsiveness of the airway, septic shock, glomerulonephritis,
inflammatory bowel disease, Crohn's disease, ulcerative colitis, atherosclerosis, growth and
metastases of malignant cells, myoblastic leukaemia, diabetes, Alzheimer's disease,
30 meningitis, osteoporosis, burn injury, ischaemic heart disease, stroke, varicose veins,

1 sarcoidosis, rhinitis, acute and chronic pain, multiple sclerosis, myeloma, bone loss
2 associated with malignancy and inflammatory and neurodegenerative diseases of the eye
3 such as scleritis, episcleritis, uveitis, Sjogrens syndrome-keratoconjunctivitis, sclerokeratitis,
4 optic neuritis, diabetic retinopathy, retinitis pigmentosa, antimalarial - induced retinopathy.
5 They are also advantageous in the treatment of infectious diseases, e.g. anthrax, in particular
6 inflammatory disease caused or exacerbated by bacterial toxins.

7
8 According to a further aspect, therefore, the present invention provides a compound of
9 formula (I), or a pharmaceutically acceptable salt, prodrug or solvate thereof, as defined
10 above for use in therapy of the human or animal body.

11
12 The invention also provides the use of a compound of formula (I), or a pharmaceutically
13 acceptable salt, prodrug or solvate thereof, as defined above, in the manufacture of a
14 medicament for use in therapy.

15
16 It will be appreciated that "therapy" also includes "prophylaxis" unless otherwise indicated.
17 The terms "therapeutic" and "therapeutically" will be understood accordingly.

18
19 In a further aspect the present invention provides a method of treating a P2X₇ receptor
20 mediated disease condition which comprises administering to a warm-blooded animal a
21 therapeutically effective amount of a compound of the formula (I) or a pharmaceutically
22 acceptable salt, prodrug or solvate thereof.

23
24 The invention further provides a method of effecting immunosuppression (e.g. in the
25 treatment of rheumatoid arthritis, osteoarthritis, irritable bowel disease, atherosclerosis or
26 psoriasis) which comprises administering a therapeutically effective amount of a compound
27 of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore
28 defined to a patient.

The invention also provides a method of treating an obstructive airways disease (e.g. asthma or COPD) which comprises administering to a patient a therapeutically effective amount of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined to a patient.

5

It will be appreciated that dosage administered will vary depending on the compound employed, the mode of administration, the treatment desired and the disorder indicated. Typically, a daily dose of active ingredient in the range of from 0.001 mg/kg to 30 mg/kg body weight is received. This daily dose may be given in divided doses as necessary, the
10 precise amount of the compound received and the route of administration depending on the weight, age and sex of the patient being treated and on the particular disease condition being treated according to principles known in the art.

The compounds of formula (I) and pharmaceutically acceptable salts, prodrug and solvates
15 thereof may be used on their own but will generally be administered in the form of a pharmaceutical composition in association with a pharmaceutically acceptable adjuvant, diluent or carrier. Depending on the mode of administration, the pharmaceutical composition will preferably comprise from 0.05 to 99 %w (per cent by weight), more preferably from 0.10 to 70 %w, of active ingredient, and, from 1 to 99.95 %w, more
20 preferably from 30 to 99.90 %w, of a pharmaceutically acceptable adjuvant, diluent or carrier, all percentages by weight being based on total composition.

The present invention therefore also provides a pharmaceutical composition comprising a compound of formula (I), or a pharmaceutically acceptable salt, prodrug or solvate thereof
25 in association with a pharmaceutically acceptable adjuvant, diluent or carrier.

The pharmaceutical compositions of the invention may be administered in standard manner for the disease condition that it is desired to treat, for example by oral, topical, parenteral, buccal, nasal, vaginal or rectal administration or by inhalation. For these purposes the
30 compounds of this invention may be formulated by means known in the art into the form of,

for example, tablets, capsules, aqueous or oily solutions, suspensions, emulsions, creams, ointments, gels, nasal sprays, suppositories, finely divided powders or aerosols for inhalation, and for parenteral use (including intravenous, intramuscular or infusion) sterile aqueous or oily solutions or suspensions or sterile emulsions.

In addition to the compounds of the present invention the pharmaceutical composition of the invention may also contain, or be co-administered (simultaneously or sequentially) with, one or more pharmacological agents of value in treating one or more disease conditions referred to above. Typically unit dosage forms will contain about 1 mg to 500 mg of a compound according to the invention.

Thus, the invention further relates to combination therapies for the treatment of any one of rheumatoid arthritis, osteoarthritis, osteoporosis, psoriasis, inflammatory bowel diseases, COPD, asthma, allergic rhinitis or cancer or the neurodegenerative diseases such as multiple sclerosis, Alzheimer's disease or stroke.

For the treatment of rheumatoid arthritis, the compounds of the invention may suitably be combined with "biological agents" such as TNF- α inhibitors such as anti-TNF monoclonal antibodies (such as Remicade, CDP-870 and Humira) and TNF receptor immunoglobulin molecules (such as Enbrel.reg.). IL-1 receptor antagonist (such as Anakinra) and IL-1 trap, IL-18 receptor, anti-IL-6 Ab, anti-CD20 Ab, anti-IL-15 Ab and CTLA4Ig.

Suitable agents to be used in combination include standard non-steroidal anti-inflammatory agents (hereinafter NSAID's) such as piroxicam, diclofenac, propionic acids such as naproxen, flubiprofen, fenoprofen, ketoprofen and ibuprofen, fenamates such as mefenamic acid, indomethacin, sulindac, apazone, pyrazolones such as phenylbutazone, salicylates such as aspirin. The COX-2 inhibitors (such as meloxicam, celecoxib, rofecoxib, valdecoxib and etoricoxib) and the cyclo-oxygenase inhibiting nitric oxide donors (CINOD's) and the "disease modifying agents" (DMARDs) such as methotrexate, sulphasalazine, cyclosporine A, lefunomide; ciclesonide; hydroxychloroquine, d-penicillamine, auranofin or parenteral or

oral gold.

The present invention also relates to the combination of a compound of the invention together with a leukotriene biosynthesis inhibitor, 5-lipoxygenase (5-LO) inhibitor or 5-lipoxygenase activating protein (FLAP) antagonist selected from the group consisting of zileuton; ABT-761; fenleuton; tepoxalin; Abbott-79175; Abbott-85761; N-(5-substituted)-thiophene-2-alkylsulfonamides; 2,6-di-tert-butylphenol hydrazones; methoxytetrahydropyrans such as Zeneca ZD-2138; the compound SB-210661; pyridinyl-substituted 2n cyanonaphthalene compounds such as L-739,010; 2-cyanoquinoline compounds such as L-746,530; indole and quinoline compounds such as MK-591, MK-886, and BAY x 1005.

The present invention still further relates to the combination of a compound of the invention together with a receptor antagonists for leukotrienes LTB₄, LTC₄, LTD₄, and LTE₄ selected from the group consisting of the phenothiazin-3-ones such as L-651,392; amidino compounds such as CGS-25019c; benzoxalamines such as ontazolast; benzenecarboximidamides such as BIIL 284/260; and compounds such as zafirlukast, ablukast, montelukast, pranlukast, verlukast (MK-679), RG-12525, Ro-245913, iralukast (CGP 45715A), and BAY x 7195.

The present invention still further relates to the combination of a compound of the invention together with a PDE4 inhibitor including inhibitors of the isoform PDE4D.

The present invention also relates to the combination of a compound of the invention together with a antihistaminic H₁ receptor antagonists including cetirizine, loratadine, desloratadine, fexofenadine, astemizole, azelastine, and chlorpheniramine.

The present invention still further relates to the combination of a compound of the invention together with a gastroprotective H₂ receptor antagonist or the proton pump inhibitors (such as omeprazole)

The present invention still further relates to the combination of a compound of the invention together with an α_1 - and α_2 -adrenoceptor agonist vasoconstrictor sympathomimetic agent, including propylhexedrine, phenylephrine, phenylpropanolamine, pseudoephedrine, naphazoline hydrochloride, oxymetazoline hydrochloride, tetrahydrozoline hydrochloride, xylometazoline hydrochloride, and ethylnorepinephrine hydrochloride.

The present invention still further relates to the combination of a compound of the invention together with anticholinergic agents including ipratropium bromide; tiotropium bromide; oxitropium bromide; pirzepine; and telenzepine.

The present invention still further relates to the combination of a compound of the invention together with a β_1 - to β_4 -adrenoceptor agonists including metaproterenol isoproterenol, isoprenaline, albuterol, salbutamol, formoterol, salmeterol, terbutaline, orciprenaline, bitolterol mesylate, and pirbuterol; or methylxanthanines including theophylline and aminophylline; sodium cromoglycate; or muscarinic receptor (M1, M2, and M3) antagonist.

The present invention still further relates to the combination of a compound of the invention together with other modulators of chemokine receptor function such as CCR1, CCR2, CCR2A, CCR2B, CCR3, CCR4, CCR5, CCR6, CCR7, CCR8, CCR9, CCR10 and CCR11 (for the C-C family); CXCR1, CXCR3, CXCR4 and CXCR5 (for the C-X-C family) and CX₃CR1 for the C-X₃-C family.

The present invention still further relates to the combination of a compound of the invention together with an insulin-like growth factor type I (IGF-1) mimetic.

The present invention still further relates to the combination of compound of the invention together with an inhaled glucocorticoid with reduced systemic side effects, including prednisone, prednisolone, flunisolide, triamcinolone acetonide, beclomethasone

dipropionate, budesonide, fluticasone propionate, and mometasone furoate.

The present invention still further relates to the combination of a compound of the invention together with (a) tryptase inhibitors; (b) platelet activating factor (PAF) antagonists; (c)
5 interleukin converting enzyme (ICE) inhibitors; (d) IMPDH inhibitors; (e) adhesion molecule inhibitors including VLA-4 antagonists; (f) cathepsins; (g) MAP kinase inhibitors; (h) glucose-6 phosphate dehydrogenase inhibitors; (i) kinin-B₁ - and B₂ -receptor antagonists; (j) anti-gout agents, e.g., colchicine; (k) xanthine oxidase inhibitors, e.g., allopurinol; (l) uricosuric agents, e.g., probenecid, sulfinpyrazone, and benzbromarone; (m)
10 growth hormone secretagogues; (n) transforming growth factor (TGF β); (o) platelet-derived growth factor (PDGF); (p) fibroblast growth factor, e.g., basic fibroblast growth factor (bFGF); (q) granulocyte macrophage colony stimulating factor (GM-CSF); (r) capsaicin cream; (s) Tachykinin NK₁ and NK₃ receptor antagonists selected from the group consisting of NKP-608C; SB-233412 (talnetant); and D-4418; and (t) elastase inhibitors selected from
15 the group consisting of UT-77 and ZD-0892 (u) induced nitric oxide synthase inhibitors (iNOS) or (v) chemoattractant receptor-homologous molecule expressed on TH2 cells, (CRTH2 antagonists).

The present invention still further relates to the combination of a compound of the invention
20 together with an inhibitor of matrix metalloproteases (MMPs), i.e., the stromelysins, the collagenases, and the gelatinases, as well as aggrecanase; especially collagenase-1 (MMP-1), collagenase-2 (MMP-8), collagenase-3 (MMP-13), stromelysin-1 (MMP-3), stromelysin-2 (MMP-10), and stromelysin-3 (MMP-11).

25 The compounds of the invention can also be used in combination with existing therapeutic agents for the treatment of osteoarthritis. Suitable agents to be used in combination include standard non-steroidal anti-inflammatory agents (hereinafter NSAID's) such as piroxicam, diclofenac, propionic acids such as naproxen, flubiprofen, fenoprofen, ketoprofen and ibuprofen, fenamates such as mefenamic acid, indomethacin, sulindac, apazone,
30 pyrazolones such as phenylbutazone, salicylates such as aspirin, induced nitric oxide

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2222
2223
2224

agonists, Dopamine agonists and inhibitors of neuronal nitric oxide synthase), and anti Alzheimer's drugs such as donepezil, tacrine, COX-2 inhibitors, propentofylline or metryfonate.


- 5 The compounds of the present invention may also be used in combination with osteoporosis agents such as roloxifene, droloxifene, lasofoxifene or fosomax and immunosuppressant agents such as FK-506, rapamycin, cyclosporine, azathioprine, and methotrexate;.

10 The activity and selectivity of the compounds according to the invention may be determined using an appropriate assay as described, for example in WO 99/29660.

The invention is further illustrated by the following non-limiting examples.
The relevant starting materials are commercially available or may be made by any convenient method as described in the literature or known to the skilled chemist or
15 described in the Examples herein.

Example 1

2-(2-Adamantyl)-N-(5-methoxy-2-methylphenyl)acetamide



Thionyl chloride (3.0 mL) and 2-adamantylacetic acid (96 mg) were heated together at 79°C for 5 minutes with stirring before cooled to room temperature and concentrated. The residue was redissolved in dichloromethane (5 mL) and added to a solution of 5-methoxy-2-methylaniline (180 mg) and triethylamine (2.0 mL) in dichloromethane (10 mL). After
25 stirring for 1 hour, dichloromethane (60 mL) and aqueous hydrochloric acid (40 mL, 2 M)

were added, the layers were separated and the organic fraction was dried, filtered and evaporated. Purification (SiO₂, dichloromethane) gave the title compound (0.12 g)


¹H NMR (400 MHz, d₆-DMSO) δ 9.15 (1H, s), 7.08 (1H, d), 7.05 (1H, d) 6.65 (1H, dd), 3.69 (3H, s), 2.49 (2H, m), 2.21 (1H, m), 2.10 (3H, s), 2.00-1.92 (2H, m), 1.89-1.67 (10H, m), 1.56-1.49 (2H, m).

MS: APCI (+ve) 314 (M+H⁺)

MP: 193-196°C

10 Example 2

2-(2-Adamantyl)-N-1H-indol-4-ylacetamide



To a solution of 1H-indol-4-amine (20 mg) in dimethylformamide (5 mL) were added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (29 mg), N,N-dimethylpyridin-4-amine (18.5 mg) and 2-adamantylacetic acid (29 mg) with continuous stirring. The mixture was heated to 60°C for 2 hours under nitrogen. The reaction was subsequently cooled to room temperature and then partitioned between ethyl acetate (20 mL) and water (20 mL). The aqueous phase was further extracted with ethyl acetate (2 x 20 mL) and the combined organics were washed with aqueous potassium hydrogen sulphate (2 x 10 mL, 2 M), saturated aqueous sodium bicarbonate (2 x 10 mL) and brine (20 mL). The organics were then dried, filtered and evaporated under vacuum to give the title compound as a white solid (23 mg).

¹H NMR (300 MHz, CDCl₃) δ 7.77-7.75 (1H, m), 7.22-7.14 (3H, m), 6.47 (1H, d), 2.60 (2H, d), 2.40 (1H, bt), 1.96-1.86 (4H, m), 1.76 (2H, s), 1.53 (6H, s).


MS: APCI (+ve) 309 (M+H⁺)

MP: 206-208°C

Example 3

2-(1-Adamantyl)-N-quinolin-5-ylacetamide

5



Prepared according to the method of example 2 using 5-quinolinamine.

MS: APCI (+ve) 321 (M+H+).

10

Pharmacological Analysis

Certain compounds such as benzoylbenzoyl adenosine triphosphate (bbATP) are known to be agonists of the P2X₇ receptor, effecting the formation of pores in the plasma membrane (Drug Development Research (1996), 37(3), p.126). Consequently, when the receptor is activated using bbATP in the presence of ethidium bromide (a fluorescent DNA probe), an increase in the fluorescence of intracellular DNA-bound ethidium bromide is observed. The increase in fluorescence can be used as a measure of P2X₇ receptor activation and therefore to quantify the effect of a compound on the P2X₇ receptor.

20

In this manner, the title compounds of the Examples were tested for antagonist activity at the P2X₇ receptor. The test was performed in 96-well flat bottomed microtitre plates, the wells being filled with 250 µl of test solution comprising 200 µl of a suspension of THP-1 cells (2.5 x 10⁶ cells/ml) containing 10⁻⁴M ethidium bromide, 25 µl of a high potassium buffer solution containing 10⁻⁵M bbATP, and 25 µl of the high potassium buffer solution containing 3 x 10⁻⁵M test compound. The plate was covered with a plastics sheet and incubated at 37 °C for one hour. The plate was then read in a Perkin-Elmer fluorescent

25

plate reader, excitation 520 nm, emission 595 nm, slit widths: Ex 15 nm, Em 20 nm. For the purposes of comparison, bbATP (a P2X₇ receptor agonist) and pyridoxal 5-phosphate (a P2X₇ receptor antagonist) were used separately in the test as controls. From the readings obtained, a pIC₅₀ figure was calculated for each test compound, this figure being the negative logarithm of the concentration of test compound necessary to reduce the bbATP agonist activity by 50%. Each of the compounds of the Examples demonstrated antagonist activity, having a pIC₅₀ figure greater than 6.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.